

Unilag Journal of Mathematics and Applications, Volume 4, Issue 1 (2024), Pages 70–80. ISSN: 2805 3966. URL: http://lagjma.edu.ng

QUASI-SYMMETRIC POLYNOMIALLY BOUNDED FRÉCHET ALGEBRAS

OLUFEMI O. OYADARE*

ABSTRACT. This paper concerns the notion of a symmetric algebra and its generalization to a quasi-symmetric algebra. We study the structure of these algebras in respect to their *hull-kernel regularity* and existence of some ideals, especially the *hull-minimal ideals*. An immediate application of our results is to the Harish-Chandra Schwartz Fréchet algebras, $C^p(G)$, of a (connected) semi-simple G.

1. INTRODUCTION

It is well-known, from J. Ludwig([8]) that every semi-simple symmetric polynomially bounded Fréchet algebra is hull-kernel regular and has a hull-minimal ideal generated by some elements of the algebra. This result gives a way of verifying the existence of hull-minimal ideals and of computing their basis elements in non-normable algebras of harmonic analysis, as has been shown for the Schwartz algebras of nilpotent and connected semi-simple groups in J. Ludwig([8]) and O. O. Oyadare ([10]) respectively.

In this paper we give a generalization of the notion of a symmetric algebra to that of a quasi-symmetric algebra and establish the importance of this generalization by showing that every semi-simple quasi-symmetric polynomially bounded Fréchet algebra is hull-kernel regular.

2. Preliminaries

For the connected semi-simple Lie group G with finite center, we denote its Lie algebra by \mathfrak{g} whose *Cartan decomposition* is given as $\mathfrak{g} = \mathfrak{t} \oplus \mathfrak{p}$. We also denote by K the analytic subgroup of G with Lie algebra \mathfrak{t} . K is then a maximal compact subgroup of G. Choose a maximal abelian subspace \mathfrak{a} of \mathfrak{p} with algebraic dual \mathfrak{a}^*

¹⁹⁹¹ Mathematics Subject Classification. Primary: 22E30. Secondary: 22E46.

Key words and phrases. Fourier Transform; Reductive Groups; Harish-Chandra's Schwartz algebras; Symmetric Polynomial Algebras; Fréchet Algebras.

^{©2024} Department of Mathematics, University of Lagos.

Submitted: August 8, 2024. Revised: October 23, 2024. Accepted: November 5, 2024.

^{*} Correspondence: femi_oya@yahoo.com.

and set $A = \exp \mathfrak{a}$. For every $\lambda \in \mathfrak{a}^*$ put

$$\mathfrak{g}_{\lambda} = \{ X \in \mathfrak{g} : [H, X] = \lambda(H)X, \forall H \in \mathfrak{a} \}$$

and call λ a restricted root of $(\mathfrak{g}, \mathfrak{a})$ whenever $\mathfrak{g}_{\lambda} \neq \{0\}$. Denote by \mathfrak{a}' the open subset of \mathfrak{a} where all restricted roots are $\neq 0$, and called the connected components the *Weyl chambers*. Let \mathfrak{a}^+ be one of the Weyl chambers, define the restricted root λ positive whenever it is positive on \mathfrak{a}^+ and denote by Δ^+ the set of all restricted positive roots. We then have the *Iwasawa decomposition* G = KAN, where N is the analytic subgroup of G corresponding to $\mathfrak{n} = \sum_{\lambda \in \Delta^+} \mathfrak{g}_{\lambda}$, and the

polar decomposition $G = K \cdot cl(A^+) \cdot K$, with $A^+ = \exp \mathfrak{a}^+$, and $cl(A^+)$ denoting the closure of A^+ .

If we set $M = \{k \in K : Ad(k)H = H, H \in \mathfrak{a}\}$ and $M' = \{k \in K : Ad(k)\mathfrak{a} \subset \mathfrak{a}\}$ and call them the *centralizer* and *normalizer* of \mathfrak{a} in K, respectively, then; (i) Mand M' are compact and have the same Lie algebra and (ii) the factor $\mathfrak{w} = M'/M$ is a finite group called the *Weylgroup*. Let $\mathfrak{a}_{\mathcal{C}}^*$ denote the complexification of \mathfrak{a}^* . \mathfrak{w} acts on $\mathfrak{a}_{\mathcal{C}}^*$ as a group of linear transformations by the requirement

$$(s\lambda)(H) = \lambda(s^{-1}H)$$

 $H \in \mathfrak{a}, s \in \mathfrak{w}, \lambda \in \mathfrak{a}_{\mathcal{C}}^*$. We then have the Bruhat decomposition

$$G = \bigsqcup_{s \in \mathfrak{w}} Bm_s B$$

where B = MAN is a closed subgroup of G and $m_s \in M'$ is the representative of s (i.e., $s = m_s M$).

Some of the most important functions on G are the *spherical functions* which we now discuss as follows. A non-zero continuous function φ on G shall be called a *(zonal) spherical function* whenever $\varphi(e) = 1$, and we have that

$$\varphi \in C(G//K) := \{ g \in C(G) : g(k_1 x k_2) = g(x), k_1, k_2 \in K, x \in G \}$$

and $f * \varphi = (f * \varphi)(e) \cdot \varphi$ for every $f \in C_c(G//K)$, where f * g denotes the convolution of f and g defined as $(f * g)(x) := \int_G f(y)g(y^{-1}x)dy$. This leads to the existence of a homomorphism $\lambda : C_c(G//K) \to \mathcal{C}$ given as $\lambda(f) = (f * \varphi)(e)$. This definition is equivalent to the satisfaction of the functional relation

$$\int_{K} \varphi(xky) dk = \varphi(x)\varphi(y), \quad x, y \in G.$$

It has been shown by Harish-Chandra ([7]) that spherical functions on G can be parametrized by members of $\mathfrak{a}_{\mathcal{C}}^*$. Indeed every spherical function on G is of the form $\varphi_{\lambda}(x) = \int_{K} e^{(i\lambda - \rho)H(xk)} dk$, $\lambda \in \mathfrak{a}_{\mathcal{C}}^*$, $\rho = \frac{1}{2} \sum_{\lambda \in \triangle^+} m_{\lambda} \cdot \lambda$, where $m_{\lambda} = dim(\mathfrak{g}_{\lambda})$,

and that $\varphi_{\lambda} = \varphi_{\mu}$ iff $\lambda = s\mu$ for some $s \in \mathfrak{w}$. Some of the well-known properties are $\varphi_{-\lambda}(x^{-1}) = \varphi_{\lambda}(x), \ \varphi_{-\lambda}(x) = \overline{\varphi}_{\overline{\lambda}}(x), \ \lambda \in \mathfrak{a}_{\mathbb{C}}^{*}, \ x \in G$, and if Ω is the *Casimir operator* on *G* then $\Omega \varphi_{\lambda} = -(\langle \lambda, \lambda \rangle + \langle \rho, \rho \rangle) \varphi_{\lambda}$, where $\lambda \in \mathfrak{a}_{\mathbb{C}}^{*}$ and $\langle \lambda, \mu \rangle := tr(adH_{\lambda} \ adH_{\mu})$ for elements $H_{\lambda}, \ H_{\mu} \in \mathfrak{a}$. The elements H_{λ} ,

 $H_{\mu} \in \mathfrak{a}$ are uniquely defined by the requirement that $\lambda(H) = tr(adH \ adH_{\lambda})$ and $\mu(H) = tr(adH \ adH_{\mu})$ for every $H \in \mathfrak{a}$. Clearly $\Omega \varphi_0 = 0$.

Let

$$\varphi_0(x) = \int_K \exp(-\rho(H(xk)))dk$$

be denoted as $\Xi(x)$ and define $\sigma: G \to \mathbb{C}$ as $\sigma(x) = ||X||$ for every $x = k \exp X \in G$, $k \in K$, $X \in \mathfrak{a}$ where $|| \cdot ||$ is a norm on the finite-dimensional space \mathfrak{a} . These two functions are spherical functions on G and there exist numbers c, d such that $1 \leq \Xi(a)e^{\rho(\log a)} \leq c(1 + \sigma(a))^d$. Also there exists $r_0 > 0$ such that $c_0 =: \int_G \Xi(x)^2(1 + \sigma(x))^{r_0} dx < \infty$. For each $0 \leq p \leq 2$ define $\mathcal{C}^p(G)$ to be the set consisting of functions f in $C^{\infty}(G)$ for which

$$||f||_{g_1,g_2;m} := \sup_G |f(g_1;x;g_2)|\Xi(x)^{-2/p}(1+\sigma(x))^m < \infty$$

where $g_1, g_2 \in \mathfrak{U}(\mathfrak{g}_{\mathbb{C}})$, the universal enveloping algebra of $\mathfrak{g}_{\mathbb{C}}, m \in \mathbb{Z}^+, x \in G$, $f(x; g_2) := \frac{d}{dt} \Big|_{t=0} f(x \cdot (\exp tg_2))$ and $f(g_1; x) := \frac{d}{dt} \Big|_{t=0} f((\exp tg_1) \cdot x)$. We call $\mathcal{C}^p(G)$ the Schwartz space on G for each $0 and note that <math>\mathcal{C}^2(G)$ is the well-known Harish-Chandra space of rapidly decreasing functions on G. The inclusions $C_c^{\infty}(G) \subset \mathcal{C}^p(G) \subset L^p(G)$ hold and with dense images. It also follows that $\mathcal{C}^p(G) \subseteq \mathcal{C}^q(G)$ whenever $0 \leq p \leq q \leq 2$. Each $\mathcal{C}^p(G)$ is closed under *involution* and the *convolution*, *. Indeed $\mathcal{C}^p(G)$ is a Fréchet algebra. We endow $\mathcal{C}^p(G/K)$ with the relative topology as a subset of $\mathcal{C}^p(G)$.

For any measurable function
$$f$$
 on G we define the spherical transform \hat{f} as
 $\hat{f}(\lambda) = \int_{G} f(x)\varphi_{-\lambda}(x)dx, \ \lambda \in \mathfrak{a}_{\mathbb{C}}^{*}$. It is known that for $f, g \in L^{1}(G)$ we have
(i) $(f * g)^{\wedge} = \hat{f} \cdot \hat{g}$ on \mathfrak{F}^{1} whenever f (or g) is right - (or left-) K -invariant;
(ii) $(f^{*})^{\wedge}(\varphi) = \overline{\hat{f}(\varphi^{*})}, \varphi \in \mathfrak{F}^{1}$; hence $(f^{*})^{\wedge} = \overline{\hat{f}}$ on \mathbb{P} : and, if we define
 $f^{\#}(g) := \int_{K \times K} f(k_{1}xk_{2})dk_{1}dk_{2}, x \in G$, then
(iii) $(f^{\#})^{\wedge} = \hat{f}$ on \mathfrak{F}^{1} .

3. Main Results

We start with fixing some notions.

3.1 Definition. A Fréchet algebra A is said to be involutive if there is a map $a \mapsto a^*$ on A such that $a^{**} = a$, $(a + b)^* = a^* + b^*$, $(\lambda a)^* = \overline{\lambda} a^*$, $\lambda \in \mathbb{C}$, and $(a \cdot b)^* = b^* \cdot a^*$, for all $a, b \in A$.

It is clear that, for each $0 , the Fréchet algebra <math>A = \mathcal{C}^p(G)$ or $\mathcal{C}^p(G//K)$ is involutive with involution $f \longmapsto f^*$ given as

$$f^*(x) = \overline{f(x^{-1})}, \ x \in G.$$

Now since the Fréchet algebra we are ultimately going to consider in this work is a Schwartz algebra we need to impose a growth condition on the members of the present abstract Fréchet algebra which fits into the general behaviour of Schwartz functions. Motivated by the estimates of members of $\mathcal{C}^p(G)$ as known above we give the following definition.

3.2 Definition. Let $A = (A, \{p_k\})$ be an involutive algebra, where $\{p_k\}$ is the defining collection of seminorms on A, and set $e(a) = \sum_{k=1}^{\infty} \frac{a^k}{k!}$ for $a \in A$. An

element $b \in A$ is said to be *polynomially bounded* if for every $k \in \mathbb{N}$ there is a constant $c_k = c_k(b) > 0$ such that

$$p_k(e(i\lambda b)) \leq c_k(1+|\lambda|)^{c_k}$$
, for all $\lambda \in \mathbb{R}$.

3.3 Remarks.

(1.) This definition may be compared with the weak inequality of the last chapter using the fact $\Xi(x) \leq 1$, for all $x \in G$.

(2.) Clearly $e(a) = \exp(a) - 1$. See J. Dixmier ([6]).

As a guiding example we consider the special (but important) case of when B is a Banach algebra, $(B, \|\cdot\|)$. In this case, if B(b) is a maximal abelian closed subalgebra of B containing b, and χ is a character on B(b) for which $\chi(b) = \mu \in spec(b)(:= spectrum of b)$, we know that in general $\mu \in \mathbb{C}$. *i.e.*, $\mu = \alpha + i\beta$ where $\alpha, \beta \in \mathbb{R}$. However we also know that, for every $\gamma \in \mathbb{R}$,

$$|e^{i\gamma\mu}| = |e^{i\gamma(\alpha+i\beta)}| = |e^{i\gamma\alpha}||e^{-\gamma\beta}| = (1)|e^{-\gamma\beta}| = e^{-\gamma\beta};$$

so that $e^{-\gamma\beta} = |e^{i\gamma\mu}| = |\exp(i\gamma\chi(b)| = |1 + e(i\gamma\chi(b))| = |1 + \chi(e(i\gamma b))| \le 1 + ||e(i\gamma b)||$. That is,

$$e^{-\gamma\beta} \le 1 + \|e(i\gamma b)\|$$

for all $b \in B$, meaning that $||e(i\gamma b)||$ grows exponentially in γ , with β a real constant.

Thus in order to have an element $b \in B = (B, |\cdot|)$ to be polynomially bounded it must be such that β in the above inequality must be zero. *i.e.*, the element bmust have real spectrum. Thus, since every Banach algebra is a Fréchet algebra we consider the requirement of having a real spectrum for a polynomially bounded element of the Fréchet algebra, $(A, \{p_k\})$. We however recall that an involutive Banach algebra in which the spectrum of every self-adjoint element is a subset of \mathbb{R} is called *symmetric* and then introduce the following notion of symmetricity.

3.4 Definition. A Fréchet algebra A is said to be *symmetric* if it admits a continuous involution and there exists a continuous *-homomorphism, σ , of A into a C^* -algebra, \mathfrak{C} , such that $spec_A(a) = spec_{\mathfrak{C}}(\sigma(a))$, for every $a \in A$ (Here $spec_A(a)$ represents the spectrum of an element a in A defined as

$$\{\lambda \in \mathbb{C} : a - \lambda \cdot 1 \text{ is not invertible}\}).$$

3.5 Remarks.

(1.) By a continuous *-homomorphism $\sigma : A \to \mathfrak{C}$ we mean a continuous homomorphism σ in which $\sigma(a^*) = \sigma(a)^{\#}$ where A and \mathfrak{C} are *- and #-involutive algebras respectively.

(2.) The requirement on the spectrum in (3.4) above is equivalent to saying that the continuous *-homomorphism, σ , is spectrum invariant.

(3.) If in (3.4) above, we require only that $spec_A(a) \subseteq spec_{\mathfrak{C}}(\sigma(a))$, then we shall refer to A a quasi-symmetric Fréchet algebra. Clearly every symmetric Fréchet algebra is automatically quasi-symmetric but not conversely. Thus the notion of quasi-symmetricity of a Fréchet algebra is more general than that of the symmetricity in (3.4). Indeed we shall in a moment extend the results of J. Ludwig ([8] Propositions 1.8 and 1.10) to include all quasi-symmetric Fréchet algebras.

We now consider the notion of a *functional calculus* (E. Hille and R. S. Phillips ([18])) that is needed shortly. We start with a Banach algebra example.

3.6 Definition. Let B be a Banach *-algebra. A function φ is said to operate on an element $f \in B$ if

- (*i.*) the Gelfand transform, \hat{f} , of f, with respect to the smallest commutative Banach sub-algebra B(f) containing f, is real, and
- (*ii.*) there exists a $g \in B(f)$ such that $\varphi \circ \hat{f} = \hat{g}$.

We shall in this case write $\varphi\{f\} = g$. Indeed, if for $f \in B$ we write $e(f) = \sum_{k=1}^{\infty} \frac{(if)^k}{k!}$ and if $||e(nf)||_B = O(||n||^N)$ as $||n|| \to \infty$, then every $\varphi \in C_c^k(\mathbb{R})$ with

k > N+1, and $\varphi(0) = 0$ operates on f and $\varphi\{f\} = \frac{1}{2\pi i} \int_{\mathbb{R}} \hat{\varphi}(\lambda) e(\lambda f) d\lambda$ so that $\|\varphi\{f\}\|_{B} \le c \|\varphi\|_{C^{k}_{c}(\mathbb{R})}.$

In a Fréchet algebra A we may also employ the functional calculus of C^{∞} functions on polynomially bounded elements of A (*cf.* J. Dixmier [6]. Let us
denote by $C^{\infty}_{c,0}(\mathbb{R})$ members $\varphi \in C^{\infty}_{c}(\mathbb{R})$ in which $\varphi(0) = 0$, then the integral $\frac{1}{2\pi i} \int_{\mathbb{R}} \hat{\varphi}(\lambda) e(i\lambda a) d\lambda$ exists and converges in A, for any polynomially bounded
elements $a \in A$ (J. Dixmier, [6] p. 18). We then define

$$\varphi\{a\} = \frac{1}{2\pi i} \int_{\mathbb{R}} \hat{\varphi}(\lambda) e(i\lambda a) d\lambda.$$

This functional calculus on polynomially bounded elements of A has the following interesting properties that makes it what we actually need.

Let a be a polynomially bounded element of the Fréchet algebra A, and let A(a) be a maximal abelian closed subalgebra containing a. If χ is any character on A(a) then $\chi(\varphi\{a\}) = \varphi\{\chi(a)\}$. We thus have

$$\chi((\psi \cdot \varphi)\{a\}) = (\psi \cdot \varphi)\{\chi(a)\} = \psi\{\chi(a)\} \cdot \varphi\{\chi(a)\} = \chi(\psi\{a\}) \cdot \chi(\varphi\{a\})$$

for $\psi, \varphi \in C^{\infty}_{c,0}(\mathbb{R})$. If A(a) is now semi-simple. That is, if $\ker(\chi) = \{0\}$ for every character χ of A(a), then

$$\chi((\psi \cdot \varphi)\{a\}) = \chi(\psi\{a\}) \cdot \chi(\varphi\{a\}) = \chi(\psi\{a\} \cdot \varphi\{a\})$$

implying $\chi((\psi \cdot \varphi)\{a\} - \psi\{a\} \cdot \varphi\{a\}) = 0$ so that

$$\psi\{a\}\cdot\varphi\{a\} = (\psi\cdot\varphi)\{a\}\cdots\cdots(I).$$

The use to which this calculus is put is contained in the following.

3.7 Proposition (J. Ludwig, [8] p. 80). Let *a* be a polynomially bounded element of *A*. Then there exist $\psi, \varphi \in C^{\infty}_{c,0}(\mathbb{R})$ such that

$$\psi\{a\}\cdot\varphi\{a\}=\varphi\{a\}.$$

Proof. Since $C_{c,0}^{\infty}(\mathbb{R})$ is (completely) regular we know that there exist $\psi, \varphi \in C_{c,0}^{\infty}(\mathbb{R})$ such that $\psi \cdot \varphi = \varphi$. Using this relation on the right-hand side of (I) above gives the result. \Box

We shall soon see how the result of Proposition 3.7 above simplifies matters in the proof of an important result of this section, thus making it central to our discussion. In an attempt to generalise this calculus to the algebra $\mathcal{C}^p(G//K)$ one may introduce a *distributional calculus* on members of $\mathcal{C}^p(G)$, 0 . SeeO. O. Oyadare ([14] and [15])

This generalisation is analogous to the generalisation of characters of finite and compact groups as functions on the groups to global characters on connected semisimple Lie groups as distributions in which J. G. Arthur ([1]), W. H. Barker ([2], [3] and [4]) and O. O. Oyadare ([9] and [11]) would prove very useful. However due to the well-developed theory of $\mathfrak{Z}(\mathfrak{g}_{\mathbb{C}})$ -finite K-finite functions and of cusp forms on connected semi-simple Lie groups no calculus or special Fourier transforms is needed in the construction of a basis for j(C) in the Schwartz algebras of focus in the last section of this chapter.

We have seen how to express a primitive Ideal of an associative algebra, A, as the kernel of some algebraically irreducible representation of A (cf. (4.2.1)). Now that we have the algebra A to be Fréchet, on which we have a topology induced by its collection of seminorms, we may employ both algebraic and topological irreducibility of a representation of A and make a comparison between them. Indeed we have the following generalisation of (1.8) in J. Ludwig ([8], p. 18) which is needed in the proof of the major result of this section.

3.8 Lemma. Every algebraically irreducible representation space of a quasisymmetric Fréchet algebra A is equivalent to a submodule of a topologically irreducible representation space of A.

Proof. Since A is quasi-symmetric so also is the adjunction $A := \mathbb{C}1 \oplus A$, of 1 to A. Thus we may assume, without any loss of generality, that A and \mathfrak{C} have identities. Now let σ be as in (3.4) and let (T, V) be an algebraically irreducible representation of A. We claim that $\ker(\sigma) \subseteq \ker(T)$.

Indeed, if $x \in \ker(\sigma)$ and $y \in A$, then the spectrum of yx in A is reduced to $\{0\}$. Suppose on the contrary to the claim, that $x \in \ker(\sigma)$ does not imply $x \in \ker(T)$, then $Tx \neq 0$; so that there exists $0 \neq v \in V$ such that $(Tx)v \neq 0$; and since T is simple, being algebraically irreducible, then we can find an element $y \in A$ such that T(y)(T(x)v) = v. *i.e.*, $(T(yx) - \lambda \cdot 1)v = 0$. This means that 1 is in the spectrum of yx, a contradiction to the fact that the spectrum of yx is reduced to $\{0\}$. Hence ker $(\sigma) \subseteq \text{ker}(T)$.

It then follows that there exists a proper maximal left Ideal M of A in which $\ker(T) \subseteq M$ in which the simple module (T, V) is equivalent to the left-regular representation of A on A/M. We note that the sum of $\mathbb{C}1$ and $\sigma(M)$ is direct in \mathfrak{C} , since otherwise $1 \in M \mod (\ker \sigma)$ which implies $1 \in \ker \sigma \subseteq (T) \subseteq M$. *i.e.*, $1 \in M$, which is impossible (see C. E. Rickart, [20], Corollary 2.1.2).

Define $M := \sigma(\mathbb{C}1 + M)$ $(= \mathbb{C}1 + \sigma(M) \subseteq \mathfrak{C})$ and define a linear functional φ on \tilde{M} by the requirement that $\varphi(\lambda \cdot 1 + \sigma(m)) = \lambda$ for every $\lambda \in \mathbb{C}$, $m \in M$. In other words for every $x \in M$, which is of the form $x = \lambda \cdot 1 + m$ we define $\varphi(\sigma(x)) = \lambda$. Now since $x = \lambda \cdot 1 + m \in M$ we have $x - \lambda \cdot 1 = m \in M$; meaning that $(x - \lambda \cdot 1)$ is non-invertible in A. *i.e.*, $\lambda \in spec_A(x)$, and by the hypothesis of quasi-symmetricity on A, we conclude that $\lambda \in Spec_{\mathfrak{C}}(\sigma(x))$. Thus $|\varphi(\sigma(x))| = |\lambda| \leq \sup\{|\mu| : \mu \in spec_{\mathfrak{C}}(\sigma(x))\} = ||\sigma(x)||_{\mathfrak{C}}$ (cf. Theorem 8 of F. F. Bonsall and J. Duncan, [16], p. 23). Therefore

$$\|\varphi\|_{op} = \sup_{\substack{x \in M \\ x \neq 0}} \frac{|\varphi(\sigma(x))|}{\|\sigma(x)\|_{\mathfrak{C}}} \le 1.$$

Hence by the Hahn-Banach theorem there exists a continuous extension, say $\tilde{\varphi}$, of φ to the whole of \mathfrak{C} of norm ≤ 1 . Since $\varphi(1) = 1$ and $\|\tilde{\varphi}\|_{op} \leq 1$, then $\tilde{\varphi}$ is a positive linear functional for which $\tilde{\varphi}(\sigma(\mathbb{C}1 + M)) = \tilde{\varphi}(\tilde{M}) = \{0\}$. Since M is maximal, we then have that $\varphi(\sigma(M)) = \{0\}$ and we can deduce that $M = \{y \in A : \tilde{\varphi}(\sigma(y^*y)) = 0\}$ which, in particular, shows that M is closed. (Indeed, every proper maximal Ideal is closed).

We therefore put a pre-Hilbert structure, $\langle \cdot, \cdot \rangle$ on A/M by setting

$$\langle x + M, y + M \rangle := \tilde{\varphi}(\sigma(y^*x)).$$

Let \mathbb{H} be the completion of $A/M = (A/M, \langle \cdot, \cdot \rangle)$, then the above left-regular representation of A on A/M extends to a unitary representation π of A on \mathbb{H} (*cf*. J. Dixmier, [17] (2.4.4)). Since we may assume that $\tilde{\varphi}$ is a *pure state* (see R. V. Kadison and J. R. Ringrose, [19], p. 213), we also know that π is topologically irreducible (*cf*. R. V. Kadison and J. R. Ringrose, [19], (2.5)). \Box

It follows from the above result that a quasi-symmetric Fréchet algebra A has sufficiently many algebraically irreducible (unitary) representations. This allows the use (2.2.9) (*i*.) of C. E. Rickart, [20], to define members of Prim(A) which leads to the discussion of the hull-minimal Ideals.

In order to then state the major result of this section we put the notion of (3.2) in the proper form in which it is needed.

3.9 Definition. An involutive Fréchet algebra A is said to be *polynomially* bounded if the set A_0 of self-adjoint polynomially bounded elements of A is dense in the real subspace A_n of hermitian elements of A.

We note here that an involutive Schwartz algebra is automatically a polynomially bounded Fréchet algebra. In particular the family of algebras given in (3.4) as $C^p(G//K)$ is a family of polynomially bounded Fréchet algebras. We now give a generalisation of (1.10) in J. Ludwig ([8], p. 81), to a semi-simple quasi-symmetric polynomially bounded Fréchet algebra. This result is the first major result of this section of the work. It assures us that there is a reasonable framework for the study of the hull-minimal Ideals in a Fréchet algebra.

3.10 Theorem. Every semi-simple quasi-symmetric polynomially bounded Fréchet algebra is hull-kernel regular.

Proof. Let A be a semi-simple quasi-symmetric polynomially bounded Fréchet algebra, then by (3.8) for every $J \in Prim(A)$ (which is of the form J = kernel of an algebraically irreducible representation of A) there exists a topologically irreducible unitary representation (π_J, \mathbb{H}_J) of A such that ker $(\pi_J) = J$.

Now let C be any closed subset of Prim(A) and fix $J \in Prim(A) \setminus C$ we need to show that there are $a_J, b_J \in A$ in which

(*i.*) $b_J \in \bigcap_{J' \in C} J'(=: \ker(C)), a_J \notin J$, and (*ii.*) $b_J \cdot a_J = a_J$.

Since C is closed there exists $u \in A$ such that

$$u \in \ker(C) \left(=\bigcap_{J' \in C} J'\right)$$

and $u \notin J$. Equivalently, $\pi_{J'}(u^*u) = 0$ for every $J' \in C$ and $\pi_J(u^*u) \neq 0$, respectively. By unitarity of $\pi_{J'}$ it is possible, after multiplying $v := u^*u$ with a positive constant, to have $\|\pi_J(v)\|_{op} = 1$. Clearly $v \in A_h$ as $v^* = v$. Also by the continuity of the *-homomorphism σ (as in (3.4)) we have that there exist a continuous seminorm p on A such that $\|\sigma(a)\|_{\mathfrak{C}} \leq p(a)$, for all $a \in A$. Thus for any unitary representation π of A we have $\|\pi(a)\|_{op} \leq \|\sigma(a)\|_{\mathfrak{C}} \leq p(a)$, $a \in A$.

As A_0 is dense in A_n we can choose $a_0 \in A_0$ such that for

$$p(a_0 - v) < 1/m, \ m = 2, 3, 4, \dots,$$

and real C^{∞} -functions φ, ψ such that ψ vanishes in a neighbourhood \mathbb{N}_{ψ} of $\left[-\frac{2}{m}, \frac{2}{m}\right], \varphi = 1$ on $\left[1 - \frac{1}{m}, 1 + \frac{1}{m}\right]$ and $\psi \cdot \varphi = \varphi$. If we now set $b_J = \psi\{a_0\}$ and $a_J = \varphi\{a_0\}$, we have, by the functional calculus on semi-simple A, that $\psi\{a_0\} \cdot \varphi\{a_0\} = \varphi\{a_0\}$ (cf. (3.7)). *i.e.*, $b_J \cdot a_J = a_J$, as required in (*ii*.) above. To now verify (*i*.) we note that

$$\begin{aligned} \|\pi_{J'}(a_0)\|_{op} &= \|\pi_{J'}(a_0 - v)\|_{op} \text{ (since } \pi_{J'}(v) = \pi_{J'}(u^*u) = 0, \text{ for every } J' \in C) \\ &\leq p(a_0 - v) \\ &\leq \frac{1}{m}, \end{aligned}$$

which shows clearly that $\pi_{J'}(a_0) \in \mathbb{N}_{\psi}$. Thus

$$\psi(\pi_{J'}(a_0)) = 0. \ i.e., \ \pi_{J'}(\psi\{a_0\}) = 0$$

implying that $\pi_{J'}(b_J) = 0$. *i.e.*, $b_J \in \ker(\pi_{J'}) = J'$. In short, $b_J \in J'$ for each $J' \in C$. *i.e.*, $b_J \in \bigcap_{J' \in C} J' = \ker(C)$, as required. Also $|||\pi_J(a_0)||_{op} - 1| = |||\pi_J(a_0)||_{op} - ||\pi_J(v)||_{op}|$ $\leq ||\pi_J(a_0) - \pi_J(v)||_{op}$ (by continuity of $|| \cdot ||_{op}$) $= ||\pi_J(a_0 - v)||_{op} \leq p(a_0 - v) \leq 1/m, \ m = 2, 3, 4, \ldots;$

implying that

$$1 - \frac{1}{m} \le \|\pi_J(a_0)\|_{op} \le 1 + \frac{1}{m}.$$

i.e., $\varphi(\pi_J(a_0)) \neq 0$. Thus

 $\pi_J(a_J) = \pi_J(\varphi\{a_0\}) = \varphi(\pi_J(a_0)) \neq 0. \ i.e., \ a_J \notin \ker(\pi_J) = J. \quad \Box$

3.11 Corollary. Let A be a semi-simple quasi-symmetric polynomially bounded Fréchet algebra and let C be a closed subset of Prim(A). Then the hull-minimal Ideal, j(C), exists and is generated by the elements a_J , $J \notin C$.

Proof. The proof follows if we combine Lemma 3.11 with Theorem 2.3 contained in U. N. Bassey and O. O. Oyadare ([5]) (or with Theorem 4.2.20 of O. O. Oyadare [10],. \Box

Lemma 3.8, Theorem 3.10 and Corollary 3.11 hold in particular for the Harish-Chandra-type algebras $A = \mathcal{C}^p(G)$ studied in O. O. Oyadare ([12] and [13]).

The last result is an improvement on (1.10) of J. Ludwig [8] as afforded by the general notion of quasi-symmetricity introduced in (4.3.5)(3.) above. We refer to U. N. Bassey and O. O. Oyadare [5], as well as O. O. Oyadare [10], for the explicit structure of the basis elements, a_J , of (type-I) hull-minimal ideals in the quasi-symmetric polynomially bounded Fréchet algebras, $C^p(G)$, of a (connected) semi-simple G.

4. Conclusion

We have been able to give a generalization of the notion of a symmetric algebra to that of a quasi-symmetric and establish the importance of this generalization by showing that every semi-simple quasi-symmetric polynomially bounded Fréchet algebra is still hull-kernel regular. An immediate application of our results is to the Harish-Chandra Schwartz Fréchet algebras, $\mathcal{C}^p(G)$, of a (connected) semisimple G.

Declarations.

Ethics Approval and consent to participate: I declare that there is no unethical conduct in the research and I also give my consent to participate in Ethics Approval. **Consent for publication:** I give my consent to the publication of this manuscript.

Availability of data and materials: Materials consulted during the research are as listed in the References below.

Competing interest: I declare that there is no competing interest in the conduct of this research.

Funding: I declare that I received no funding for this research.

Authors' contributions: I declare that I was the only one who contributed to the conduct of this research.

Acknowledgments: I acknowledge authors whose research works were consulted as listed in the References below.

References

- Arthur, J. G. Some tempered distributions on semi-simple groups of real rank one. Ann. of Math. 100 (1974): 553-584.
- Barker, W. H. The spherical Bochner theorem on semi-simple Lie groups. J. Funct. Anal. 20 (1975): 179-207.
- [3] Barker, W. H. Positive definite distributions on unimodular Lie groups. Duke Math. J. 43. 1 (1976): 71-79.
- Barker, W. H. Tempered, invariant, positive-definite distributions on SU(1,1)/{±1}. Illinois J. Math. 28. 1 (1984): 83-102.
- [5] Bassey, U. N. and Oyadare, O. O. On Hull-minimal ideals in the Schwartz algebras of spherical functions on a semi-simple Lie group. International Journal of Functional Analysis, Operator Theory and Applications. 3 (1) (2011): 37-52.
- [6] Dixmier J. Opérateurs de rang fini dans les représentations unitaires. Inst. Hautes Études. Sci. Publ. Math. 6 (1960): 13-25.
- [7] Harish-Chandra Spherical functions on a semi-simple Lie group, I. Amer. J. Math. 80. 2 (1958): 241-310.
- [8] Ludwig, J. Hull-minimal Ideals in the Schwartz algebra of the Heisenberg group. Studia Mathematica. 130, 1 (1998), pp. 77-98.
- [9] Oyadare, O. O. On harmonic analysis of spherical convolutions on semi-simple Lie groups. Theoretical Mathematics and Applications, 5 (3) (2015), p. 19-36.
- [10] Oyadare, O. O. On ideal theory for the Schwartz algebras of spherical functions on semisimple Lie groups. Ph.D Thesis, University of Ibadan (2016). x + 383 pp.
- [11] Oyadare, O. O. Functional analysis of canonical wave-packets on real reductive groups. arXiv:1912.07542v1. [math.FA], 13 Dec. 2019.
- [12] Oyadare, O. O. Non-spherical Harish-Chandra Fourier transforms on real reductive groups. J. Fourier Anal. Appl.. 28, 15 (2022). http://doi.org/10.1007/s00041-09906-w
- [13] Oyadare, O. O. The full Bochner theorem on real reductive groups. Algebras, Groups and Geometries 39, (2023), p. 207-220.
- [14] Oyadare, O. O. Series analysis and Schwartz algebras of spherical convolutions on semisimple Lie groups. Algebras, Groups and Geometries. 40(1), (2024a), p. 41-59. See also arXiv:1706.09045 [math.RT].
- [15] Oyadare, O. O. On the operator-valued Fourier transform of the Harish-Chandra Schwartz Algebra. arXiv:2407.20755v1. [math.RT], 2024b.

OLUFEMI O. OYADARE

- [16] Bonsall, F. F. and Duncan, J. Complete normed algebras. Berlin: Springer-Verlag, 1973.
- [17] Dixmier, J. Les C*-algebres et leurs representations. Paris: Gauthier-Villars, 1964.
- [18] Hille, E. and Phillips, R. S. Functional analysis and semi-groups. 31. Rhode Island: Amer. Math. Soc., Colloq. Publ., Providence, 1957.
- [19] Kadison, R. V. and Ringrose, J. R. Fundamentals of the theory of operators algebras, vol.
 1, Elementary theory. London: Academic Press, 1983.
- [20] Rickart, C. E. General theory of Banach algebras. New York: Van Nostrand, 1974.

Olufemi O. Oyadare*

DEPARTMENT OF MATHEMATICS,, OBAFEMI AWOLOWO UNIVERSITY,, ILE-IFE, 220005, NIGE-RIA.,

E-mail address: femi_oya@yahoo.com